

# ECON 441

## Introduction to Mathematical Economics

Div Bhagia

Lecture 1: Preliminaries

# Today's Topics & References

- Numbers and sets (sections 2.2 and 2.3)
- Relations and functions (sections 2.4-2.6, page 163)
- Summation notation (handout)
- Necessary and sufficient conditions (beginning of 5.1)

# Numbers and Sets

# Real-Number System

- *Integers:*

..., -3, -2, -1, 0, 1, 2, 3, ...

# Real-Number System

- *Integers:*

..., -3, -2, -1, 0, 1, 2, 3, ...

- *Fractions:*

$\frac{1}{2}, \frac{3}{5}, -\frac{2}{3}$

# Real-Number System

- *Integers:*

..., -3, -2, -1, 0, 1, 2, 3, ...

- *Fractions:*

$\frac{1}{2}, \frac{3}{5}, -\frac{2}{3}$

- *Rational numbers:* ratio of integers

# Real-Number System

- *Integers:*

..., -3, -2, -1, 0, 1, 2, 3, ...

- *Fractions:*

$\frac{1}{2}$ ,  $\frac{3}{5}$ ,  $-\frac{2}{3}$

- *Rational numbers:* ratio of integers

*Are fractions rational numbers? What about integers?*

# Real-Number System

- *Rational numbers*: ratio of integers  
“terminating or repeating decimal”

Example.  $\frac{1}{3} = 0.333$ ,  $\frac{1}{4} = 0.25$



# Real-Number System

- *Rational numbers*: ratio of integers  
“terminating or repeating decimal”

Example.  $\frac{1}{3} = 0.333$ ,  $\frac{1}{4} = 0.25$

- *Irrational numbers*: cannot be expressed as a ratio of two integers  
“nonterminating or nonrepeating decimal”

Example.  $\sqrt{2} = 1.4142$ ,  $\pi = 3.1415$

# Real-Number System

- *Rational numbers*: ratio of integers  
“terminating or repeating decimal”

Example.  $\frac{1}{3} = 0.333$ ,  $\frac{1}{4} = 0.25$

- *Irrational numbers*: cannot be expressed as a ratio of two integers  
“nonterminating or nonrepeating decimal”

Example.  $\sqrt{2} = 1.4142$ ,  $\pi = 3.1415$

- *Real numbers* ( $\mathbb{R}$ ): *rational and irrational*

# Sets

- A set is a collection of distinct objects.

$$A = \{brownies, icecream, pizza, ramen\}$$

- $pizza \in A$ ,  $\in$  stands for 'is in'

# Sets

- A set is a collection of distinct objects.

$$A = \{brownies, icecream, pizza, ramen\}$$

- $pizza \in A$ ,  $\in$  stands for 'is in'
- What about sets  $B$  and  $C$ ?

$$B = \{x | x \text{ is a positive integer}\}$$

$$C = \{x | 1 < x < 5\}$$

# Set Relations

## 1. Equivalence (=)

$$A = \{brownies, icecream, pizza, ramen\}$$

$$B = \{pizza, ramen, icecream, brownies\}$$

$$A = B$$

# Set Relations

## 1. Equivalence (=)

$$A = \{brownies, icecream, pizza, ramen\}$$

$$B = \{pizza, ramen, icecream, brownies\}$$

$$A = B$$

## 2. Subset ( $\subset$ )

$$C = \{pizza, ramen\}$$

$C \neq A$  but  $C \subset A$ .

Note:  $A \supset C$  is equivalent.

# Set Relations

## 1. Equivalence (=)

$$A = \{brownies, icecream, pizza, ramen\}$$

$$B = \{pizza, ramen, icecream, brownies\}$$

$$A = B$$

## 2. Subset ( $\subset$ )

$$C = \{pizza, ramen\}$$

$C \neq A$  but  $C \subset A$ .

Note:  $A \supset C$  is equivalent.

Is  $A \subset B$ ?

# Set Relations

## 1. Equivalence (=)

$$A = \{brownies, icecream, pizza, ramen\}$$

$$B = \{pizza, ramen, icecream, brownies\}$$

$$A = B$$

## 2. Subset ( $\subset$ )

$$C = \{pizza, ramen\}$$

$C \neq A$  but  $C \subset A$ .

Note:  $A \supset C$  is equivalent.

Is  $A \subset B$ ? Yes, but  $C$  is a proper subset of  $A$ .



# Set Relations

## 3. Disjoint sets

$$A = \{brownies, icecream, pizza, ramen\}$$

$$D = \{salad, fruits\}$$

# Set Relations

## 3. Disjoint sets

$$A = \{brownies, icecream, pizza, ramen\}$$

$$D = \{salad, fruits\}$$

## 4. Neither but still related

$$A = \{brownies, icecream, pizza, ramen\}$$

$$E = \{salad, fruits, icecream\}$$

# Set Relations

- $\emptyset$ : empty or null set

# Set Relations

- $\emptyset$ : empty or null set
- What are all possible subsets of

$$S = \{a, b, c\}$$

# Set Relations

- $\emptyset$ : empty or null set
- What are all possible subsets of

$$S = \{a, b, c\}$$

$$\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}$$

# Set Relations

- $\emptyset$ : empty or null set
- What are all possible subsets of

$$S = \{a, b, c\}$$

$\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}$

Always  $2^n$  subsets. Here  $n = 3$ , so 8 subsets.

# Set Operations

1. *Union*:  $A \cup B$ , elements in either  $A$  or  $B$
2. *Intersection*:  $A \cap B$ , elements in both  $A$  and  $B$

Example:

$$A = \{brownies, icecream, pizza, ramen\}$$

$$B = \{salad, fruits, icecream\}$$

$$A \cup B =$$

$$A \cap B =$$

# Set Operations

1. *Union*:  $A \cup B$ , elements in either  $A$  or  $B$
2. *Intersection*:  $A \cap B$ , elements in both  $A$  and  $B$

What about

$$A = \{brownies, icecream, pizza, ramen\}$$

$$B = \{salad, fruits\}$$

$$A \cup B =$$

$$A \cap B =$$



# Set Operations



3. *Complement of A:  $\tilde{A}$ , 'not A'*

Universal set  $U$  (context specific) then:

$$\tilde{A} = \{x \mid x \in U \text{ and } x \notin A\}$$

Example.  $U = \{1, e, f, 2\}$ ,  $A = \{1, 2\}$ , then  $\tilde{A} = \{e, f\}$ .

# Set Operations: Venn Diagrams



# Laws of Set Operations

- Commutative law

$$A \cup B = B \cup A \quad A \cap B = B \cap A$$

- Distributive law


$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$


# Distributive law

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$A$




$B \cap C$




# Distributive law

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$A \cup B$



$A \cup C$



# Ordered Sets

- We said order does not matter for sets
- But we can have ordered sets where

$$(a, b) \neq (b, a) \text{ unless } a = b$$

- Ordered pairs, triples,...

Example. (*age, weight*), (22, 120) different from (120, 22)

# Cartesian Product


$$A = \{1, 2\} \quad B = \{3, 4\}$$

Cartesian Product: set of all possible ordered pairs

$$A \times B = \{(1, 3), (1, 4), (2, 3), (2, 4)\}$$

# Cartesian Plane

$$\mathbb{R}^2 = \{(x, y) \mid x \in \mathbb{R}, y \in \mathbb{R}\}$$



Can have  $\mathbb{R}^3, \mathbb{R}^4, \dots, \mathbb{R}^n$




# Relations and Functions

# Relations

Relation: subset of the Cartesian product

Example.  $\{(x, y) \mid y \leq x\}$



# Functions

Function: a relation where for each  $x$  there is a unique  $y$

$$f : X \rightarrow Y, \quad y = f(x)$$

Examples.  $y = x, y = x^2, y = 2x + 3$

$X$  : domain,  $Y$  : codomain,  $f(X)$  : range

Most functions we will encounter,  $f : \mathbb{R}^k \rightarrow \mathbb{R}$

# Functions

Let's say,

$$f : X \rightarrow \mathbb{R}, \quad y = 3x - 5$$

where  $X = \{2, 3, 4\}$ .

What is the range?

# Cost Function

Consider the total cost  $C$  of producing hats  $Q$ ,

$$C = f(Q)$$

- Should I be able to produce 10 hats and 5 hats at the same cost?
- Possible to produce 5 hats for \$20 and \$25?

## By the way

Consider the total cost  $C$  of producing hats  $Q$ ,

$$C = f(Q) = 2Q + 5$$

What is the cost of producing 1 hat?

What is the cost of producing 2 hats?

How many hats can I produce for \$25?

# Types of Functions

- Constant:  $y = f(x) = 5$
- Polynomial of degree  $n$ 
  - $n = 0$ , constant
  - $n = 1$ , linear
  - $n = 2$ , quadratic
  - $n = 3$ , cubic
  - ...
- Rational function: ratio of two polynomial functions:

$$y = \frac{a}{x}$$

# Function of More than One Variables

Functions can be of two variables:

$$z = g(x, y)$$

Or three, or four,..., or  $n$



# Monotonic functions

Strictly increasing function:

$$x_1 > x_2 \rightarrow f(x_1) > f(x_2)$$

Strictly decreasing function:

$$x_1 > x_2 \rightarrow f(x_1) < f(x_2)$$

Increasing function:

$$x_1 > x_2 \rightarrow f(x_1) \geq f(x_2)$$

Decreasing function:

$$x_1 > x_2 \rightarrow f(x_1) \leq f(x_2)$$

# Inverse of a function

Function  $y = f(x)$  has an inverse if it is a one-to-one mapping, i.e. each value of  $y$  is associated with a unique value of  $x$ .

Inverse function

$$x = f^{-1}(y)$$

returns the value corresponding value of  $x$  for each  $y$ .

One-to-one mapping unique to strictly monotonic functions

# Inverse of a function

Example: Find the inverse of  $y = f(x) = 3x - 2$ .

## By the way

What is  $x \times x$ ?

What is  $x^2 \times x$ ?

What is  $x^2 \times x^2$ ?

More generally,  $x^n \times x^m = x^{m+n}$

# Summation Notation

# Summation Notation

$$\sum_{i=1}^N x_i = x_1 + x_2 + \dots + x_N$$

Example:

$$x = \{2, 9, 6, 8, 11, 14\}$$

$$\sum_{i=1}^4 x_i = x_1 + x_2 + x_3 + x_4 = 2 + 9 + 6 + 8 = 25$$

# Summation Notation

Another way of using a summation sign is to write

$$\sum_{x \in A} x$$

which refers to summing up all elements in  $A$ .

To sum up  $x$  for all possible values  $x$ , we can simply write

$$\sum_x x$$

# Things you CAN do

1. Pull constants out of or into the summation sign.

$$\sum_{i=1}^N bx_i = b \sum_{i=1}^N x_i$$



# Things you CAN do

2. Split apart (or combine) sums (addition) or differences (subtraction)

$$\sum_{i=1}^N (bx_i + cy_i) = b \sum_{i=1}^N x_i + c \sum_{i=1}^N y_i$$

# Things you CAN do

3. Multiply through constants by the number of terms in the summation

$$\sum_{i=1}^N (a + bx_i) = aN + b \sum_{i=1}^N x_i$$

# Things you CANNOT do

1. Split apart (or combine) products (multiplication) or quotients (division).

$$\sum_{i=1}^N x_i y_i \neq \sum_{i=1}^N x_i \times \sum_{i=1}^N y_i$$

# Things you CANNOT do

2. Move the exponent out of or into the summation.

$$\sum_{i=1}^N x_i^a \neq \left( \sum_{i=1}^N x_i \right)^a$$

# Necessary and Sufficient Conditions

# Necessary vs. Sufficient Conditions

$q$  is a necessary condition for  $p$  if:

$$p \implies q$$

# Necessary vs. Sufficient Conditions

$q$  is a necessary condition for  $p$  if:

$$p \implies q$$

$p$ : I ate tofu for dinner

$q$ : My dinner had protein

# Necessary vs. Sufficient Conditions

$q$  is a sufficient condition for  $p$  if:

$$p \longleftarrow q$$



# Necessary vs. Sufficient Conditions

$q$  is a sufficient condition for  $p$  if:

$$p \longleftarrow q$$

$p$ : A number is even

$q$ : A number is divisible by 4

# Necessary vs. Sufficient Conditions

$q$  is both necessary and sufficient for  $p$

$$p \iff q$$

# Necessary vs. Sufficient Conditions

$q$  is both necessary and sufficient for  $p$

$$p \iff q$$

$p$ : A number is even

$q$ : A number is divisible by 2

# Necessary vs. Sufficient Conditions

$p$ : It is a holiday

$q$ : It is Thanksgiving

# Necessary vs. Sufficient Conditions

$p$ : The car is out of gas

$q$ : The car isn't starting

# Necessary vs. Sufficient Conditions

$p$ : A geometric figure has four sides

$q$ : It is a rectangle

# Homework Questions

- Exercise 2.3: 1, 2
- Exercise 2.4: 5, 7, 8
- Exercise 2.5: 1 (For each part, find the inverse of the function too.)
- Exercise 4.2: 6, 8
- Exercise 5.1: 1